ELSEVIER

Journal of Organometallic Chemistry 503 (1995) C57–C58

Journal of Organometallic Chemistry

Author Index of Volume 503

Abad, M.M., 193
Alexander, C.W., 213
Alonso, F.J.G., 59
Arif, A.M., 235
Ariza-Castolo, A., C40
Arndt, P., 221
Atwood, D.A., C4

Baker, P.K., C8
Baumann, W., 221
Baumeister, U., 93
Behrens, U., 101
Ben-David, Y., 149
Benito, A., 259
Biersack, M., 277
Blaschette, A., 155
Burlakov, V.V., 221

Calogero, S., 297
Canty, A.J., C16
Carriedo, G.A., 59
Casarin, M., 243
Cecchi, P., 297
Chen, W., 47
Chen, Z., 53
Cheung, K.-K., C19
Chiarini, M., 297
Chiusoli, G.P., 21
Churakov, A.V., C39
Claramunt, R.M., 265
Clinet, J.C., C48
Coles, S.J., C8
Contreras, R., C40
Corey, J.Y., 205
Costa, M., 21
Cucciolito, M.E., 251
Cunningham, D., C46

De Felice, V., 251
De Jesús Rosales-Hoz, M., C40
Dieter, R.K., 213
Domínguez-Durán, M.E., C40
Duñach, E., C48

Edwards, A.J., 15, C29
Eisen, M.S., 307
Elguero, J., 265
Ernst, R.D., 29
Eujen, R., C51

Ferguson, G., C12
Fernández-Catuxo, L., 59
Floriani, C., 185
Fröhlich, H.-O., 135

Gabriele, B., 21
Gallagher, J.F., C12
George, T.A., C1
Gervasio, G., 185
Gioia Lobbia, G., 297
Giordano, F., 251
Gladysz, J.A., 235
Glaser, M., C32
Goh, L.Y., 47
Gómez-Elipe, P., 59
Görls, H., 135
Grove, D.M., C26
Gspaltl, P., 129
Guennec, N., 171
Győryová, K., 1

Hahn, F.E., C32
Halterman, R.L., 53
Hammund, H.H., C1
Hassler, K., 93
Hayashi, Y., 143
Hengge, E., 129
Henschel, D., 155
Herberhold, M., 243, 277
Herskovics-Korine, D., 307
Hoge, B., C51
Holloway, C.E., 1
Horchler von Locquenghien, K., 289
Huang, X., 11
Hursthouse, M.B., C8

Imhof, W., 69

Jegier, J.A., C4
Jin, G.-X., 243
Jin, H., C16
Jones, P.G., 155
Juang, J.-P., 117

Kaneta, N., 143
Kapteijn, G.M., C26
Kazennova, N.B., C39
Kempe, R., 221
Kennedy, J.D., C12
Kivekäs, R., 193
Krahl, J., 155
Kundler, S., 289
Kuz'mina, L.G., C39

Lang, H., 69
Lange, I., 155
Leadbeater, N.E., 15
Lee, K.K.H., C43
Lee, S.-M., C19
Lefeber, C., 221
Leise, M., 69
Lemenovskii, D.A., C39
Lewis, J., 15
Liao, F.-L., 35
Liaw, J.-H., 35
Lin, S.-Y., 213
Lin, T.-S., 35
López, C., 265
Lu, Y.-J., 35
Lugan, N., C22
Lügger, T., C32

Ma, H., 11
Main, L., 75
Manning, A.R., C46
Martin, K.J., C4
Martínez-Máñez, R., 259
Mathieu, R., C22
McArdle, P.A., C46
Meehan, M.M., C8
Melnik, M., 1
Milstein, D., 149
Moinet, C., 171

Nicholson, B.K., 75
Nombel, P., C22
Nuñez, R., 193

O'Connell, D., C12
O'Dwyer, L., C46

Pardi, L., 243
Paver, M.A., C29
Payá, J., 259
Paz-Sandoval, M.A., C40
Pazos-Mayen, C., C40
Pearson, P., C29
Peng, T.-S., 235
Piarulli, U., 185
Portnoy, M., 149

Raithby, P.R., 15, C29
Rath, N.P., 205
Rennie, M.-A., C29
Rheingold, A.L., 29

Ridyard, S.D., C8
Rosenthal, U., 221
Rossell, O., 225
Rufanov, K.A., C39
Ruffo, F., 251
Russell, C.A., C29
Rutherford, D., C4

Sakai, H., 143
Salerno, G., 21
Sanaú, M., 235
Sanz, D., 265
Schenzel, K., 93
Schmidt, G., 101
Seco, M., 225
Segalés, G., 225
Shaltout, R.M., 205
Sillanpää, R., 193
Sinn, E., 259
Skelton, B.W., C16
Skoršepa, J., 1
Soto, J., 259
Spalding, T.R., C12
Spielberger, A., 129
Spies, H., C32
Stievano, L., 297

Teixidor, F., 193
Tendero, M.J.L., 259
Tillack, A., 221
Töllner, K., 111
Trofimenko, S., 265
Tully, W., 75

Uemura, M., 143

Valle, G., 297
Van den Broek, M.A.F.H., C26
Van Koten, G., C26
Verhoef, M.J., C26
Viñas, C., 193
Viterbo, D., 185

Wadepohl, H., 111
Waldman, T.E., 29
Wang, S.-L., 35
Wang, T.-F., 117
Wang, Y.-P., 35
Weinmann, M., 69
Wen, Y.-S., 117
White, A.H., C16
Williams, D.N., 185
Winter, M., 69
Wirth, A., 155
Wong, R.C.S., 47
Wong, W.-T., C19
Wong, W.T., C43
Wrackmeyer, B., 289
Wright, D.S., C29
Wu, Z., 11
Wyrwa, R., 135

Xu, Z., 11

Yang, Y.-H., 35
You, X., 11

Zanello, P., 243
Zechmann, A., 129
Zhou, X., 11

ELSEVIER

Journal of Organometallic Chemistry 503 (1995) C59–C68

Journal of Organometallic Chemistry

Subject Index of Volume 503

Acetylene complex
Protonation of $Cp(CO)(PPh_3)RuC{\equiv}CPh$: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct $[Cp(CO)(PPh_3)Ru{=}\overline{CCH(Ph)CH_2CH_2O}]^+$ (P. Nombel, N. Lugan and R. Mathieu), C22

Acyl
Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26

Adduct
Coordination of the (μ-η^2-As_2) ligand to Group 6 transition metal carbonyl fragments and crystal structures of $Cp_2Cr_2(CO)_4As_2$-$[M(CO)_5]_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47

Alkene
Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β-unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Alkenes
Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of [{Pt(2,9-dimethyl-1,10-phenanthroline)- (Z-$MeO_2CCH{=}CHCO_2$-$Me)(H_2O)$}$_2$ $Hg](BF_4)_2$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251

Alkyl
Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117

Alkyl chloride
Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C–Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149

Alkyne
Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)-β-diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal–organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101

Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117

Alkynes
Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21

Allylic alkylation
New chiral chelating phosphine complexes containing tricarbonyl(η^6-arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Aluminium
Tetradentate -N_2O_2 ligand complexes of Tin(II). X-ray crystal structure of [*N,N'*-(1,2-ethylene)*bis*(salicylaldamine)]tin(II), ($SaleanH_2Sn$) (D.A. Atwood, J.A. Jegier, K.J. Martin and D. Rutherford), C4

α-Aminoalkyl cuprate conjugate addition
Conjugate addition reactions of α-azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

Antimony
Synthesis and structure of the dimeric organo-antimony (V) imido complex [$Ph_3Sb(\mu$-$NCH_2CH_2Ph)]_2$ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29

η^6-Arene complex
Synthesis and chacterization of low valent complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1

Arene complexes
Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiényl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171

Arsenic
Coordination of the (μ-η^2-As_2) ligand to Group 6 transition metal carbonyl fragments and crystal structures of $Cp_2Cr_2(CO)_4As_2$-$[M(CO)_5]_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47

Donor-free bis(trifluoromethyl)cadmium, $(CF_3)_2Cd$: a readily available low-temperature difluorocarbene source (R. Eujen and B. Hoge), C51

Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplexkationen: Synthese und Strukturen von $[R_3Sn(L)(L')]^+[(MeSO_2)_2N]^-$ (L = L' = HMPA, Ph_3AsO, Imidazol, 4-Dimethylaminopyridin; L = H_2O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155

Azepine complexes
Cyclopentadienylcobalt complexes of some azepines (H. Wadepohl and K. Töllner), 111

α-Azoalkyl anion
Conjugate addition reactions of α-azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

α-Azoalkyl cuprate
Conjugate addition reactions of α-azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

Benzamidinate
Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307

η^6-Benzene complex
Synthesis and chacterization of low valent complexes of molybdenum containing the tridentate ligand PhP($CH_2CH_2PPh_2$)$_2$ (TRI) (T.A. George and H.H. Hammund), C1

Borates
Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289

Boron
Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of [$SnCl_3$(HB(4-Cl-3,5-$Me_2Pz)_3$)] and [$PhSnCl_2$(HB(4-Cl-3,5-Me_2Pz)$_3$)] (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297
Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289

2-Butyne
Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, [WI_2(CO)(η^2-triphos)(η^2-MeC$_2$-Me)].0.75CH_2Cl_2 {triphos = PhP($CH_2CH_2PPh_2$)$_2$} (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8

Cadmium
Donor-free bis(trifluoromethyl)cadmium, $(CF_3)_2Cd$: a readily available low-temperature difluorocarbene source (R. Eujen and B. Hoge), C51

Carbene complex
Protonation of Cp(CO)(PPh_3)RuC≡CPh: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct [Cp(CO)(PPh_3)Ru=CCH(Ph)CH_2CH_2O]$^+$ (P. Nombel, N. Lugan and R. Mathieu), C22

Carbonyl
Coordination of the (μ-η^2-As_2) ligand to Group 6 transition metal carbonyl fragments and crystal structures of $Cp_2Cr_2(CO)_4As_2$-[$M(CO)_5$]$_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47
Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-*S*-benzoylsulphenamide. Crystal and molecular structures of [$Os_6(CO)_{20}$(μ_4-S)(MeCN)] (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19
Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4-nitrene ligand. Crystal structures of [$Ru_6(CO)_{15}$(μ-CO)$_2$(μ_4-NH)(μ-OMe){μ_3-η^2-N(H)C(O)OMe}] and [$Ru_6(CO)_{16}$(μ-CO)$_2$(μ_4-NH)(μ-OMe)(μ-NCO)] (K.K.H. Lee and W.T. Wong), C43
The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15

μ_3-Carbonyl
The cleavage of a coordinated SCNR group in [$Fe(CO)_2L_2$(η^2-SCNR)] by [Co(η-C_5H_5)(PPh_3)$_2$] to give complexes containing μ_3-CNR ligand. The preparation and structure of [{Co(η-C_5H_5)}$_2$(Fe(CO)$_2$(PPh_3))(μ_3-S)(μ_3-CNC(O)C_6H_5)] (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46

Carbonylation
Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26

Carbonyl complexes
Formation of a rhenium(III) carbonyl complex by electrophilic attack on rhenium isocyanides. Synthesis and molecular structure of {Re[N(CH_2CH_2S)$_3$][CNC(CH_3)$_3$]} and {Re[N(CH_2CH_2S)$_3$](CO)} (M. Glaser, H. Spies, T. Lügger and F.E. Hahn), C32

Carbonyls
Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59

Carborane
Procedure for the degradation of 1,2-(PR_2)$_2$-1,2-dicarba-*closo*-dodecaborane(12) and 1-(PR_2)-2-R'-1,2-dicarba-*closo*-dodecaborane(12) (F. Teixidor, C. Viñas, M.M. Abad, R. Nuñez, R. Kivekäs and R. Sillanpää), 193
The relationship between the molecular structure of [3,3-(PMe_2Ph)$_2$-*closo*-3,1,2-$PtC_2B_9H_{11}$] and the mechanism of the fluxional behaviour of M(PR_3)$_2$-units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12

Catalysis
Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21
New chiral chelating phosphine complexes containing tricarbonyl(η^6-arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Cationic
Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59

C–C coupling
Intramolekulare Insertion eines η^5-Cyclopentadienyl-Ringes in einem Bis-η^5-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefeber, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221

Chalcones
Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β-unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Chelate
Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C–Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149

Chiral
Crystal structure of enantiomerically pure bis(dimethylbicyclooctylcyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

Chiral ligands
Transition metal carbohydrate chemistry. Part 4. Homoleptic diacetoneglucose complexes of molybdenum and tungsten: chiral ligands associated with an M≡M triple bond (U. Piarulli, D.N. Williams, C. Floriani, G. Gervasio and D. Viterbo), 185

Chiral metal–metal complexes
Transition metal carbohydrate chemistry. Part 4. Homoleptic diacetoneglucose complexes of molybdenum and tungsten: chiral ligands associated with an M≡M triple bond (U. Piarulli, D.N. Williams, C. Floriani, G. Gervasio and D. Viterbo), 185

Chiral phosphine ligand
New chiral chelating phosphine complexes containing tricarbonyl(η^6-arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Chiral recognition
Crystal structures of diastereomeric styrene complexes of the chiral rhenium fragment [(η^5-C_5H_5)Re(NO)(PPh_3)]$^+$: effect of ligating enantioface upon bonding and conformation (M. Sanaú, T.-S. Peng, A.M. Arif and J.A. Gladysz), 235

Chromium
- Coordination of the (μ-η^2-As_2) ligand to Group 6 transition metal carbonyl fragments and crystal structures of $Cp_2Cr_2(CO)_4As_2[M(CO)_5]_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47
- Syntheses and spectra of $(CO)_2(NO)Cr(\eta^5\text{-}C_5H_4\text{-}R)$ (R = –C(O)–N_3, –N=C=O, –NH_2, –NH–C(O)–O-$CH_2C_6H_5$ or –NH–C(O)–NH–$(\eta^5\text{-}C_5H_4)Cr(CO)_2(NO)$) and crystal structure of $(CO)_2(NO)Cr[(\eta^5\text{-}C_5H_4)$–NH–C(O)–NH–$(\eta^5\text{-}C_5H_4)]Cr(CO)_2$-(NO) (Y.-P. Wang, T.-S. Lin, Y.-H. Yang, J.-H. Liaw, S.-L. Wang, F.-L. Liao and Y.-J. Lu), 35

(Arene)chromium
- New chiral chelating phosphine complexes containing tricarbonyl(η^6-arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Cluster
- Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-*S*-benzoylsulphenamide. Crystal and molecular structures of $[Os_6(CO)_{20}(\mu_4\text{-}S)(MeCN)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19

Clusters
- Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic $[Mn_3(CO)_{12}(\mu\text{-}H)]^{2-}$ and $[Fe_3(CO)_{11}]^{2-}$ anions (O. Rossell, M. Seco and G. Segalés), 225
- Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4-nitrene ligand. Crystal structures of $[Ru_6(CO)_{15}(\mu\text{-}CO)_2(\mu_4\text{-}NH)(\mu\text{-}OMe)(\mu_3\text{-}\eta^2\text{-}N(H)C(O)OMe)]$ and $[Ru_6(CO)_{16}(\mu\text{-}CO)_2(\mu_4\text{-}NH)(\mu\text{-}OMe)(\mu\text{-}NCO)]$ (K.K.H. Lee and W.T. Wong), C43

^{13}C NMR
- An ^{1}H and ^{13}C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ^{13}C–^{11}B and ^{13}C–^{205}Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265

Cobalt
- Cyclopentadienylcobalt complexes of some azepines (H. Wadepohl and K. Töllner), 111
- The cleavage of a coordinated SCNR group in $[Fe(CO)_2L_2(\eta^2\text{-}SCNR)]$ by $[Co(\eta\text{-}C_5H_5)(PPh_3)_2]$ to give complexes containing μ_3-CNR ligand. The preparation and structure of $[\{Co(\eta\text{-}C_5H_5)\}_2\{Fe(CO)_2(PPh_3)\}(\mu_3\text{-}S)(\mu_3\text{-}CNC(O)C_6H_5)]$ (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46

Complexes
- Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)-β-diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal–organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101
- Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59

Copper
- Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)-β-diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal–organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101
- Reaktionsverhalten Alkinyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69

Crystal structure
- Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307
- Crystal structure of enantiomerically pure bis(dimethylbicyclooctylcyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53
- Hydrogen-bonding 'Pd(OH)...2(HOAr)' and '$[Pd(OH_2)^+ \ldots {}^-OAr]_2$', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts $[Pd(OH)(C_4H_8)\{(pz)_3BH\}.2(3\text{-}MeC_6H_4OH)]$ and $[Pd(OH_2)(C_4H_8)\{(pz)_3BH\}.(C_6F_5O)]_2$ [$(pz)_3BH$ = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16
- Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-*S*-benzoylsulphenamide. Crystal and molecular structures of $[Os_6(CO)_{20}(\mu_4\text{-}S)(MeCN)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19
- Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)-β-diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal–organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101
- Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β-unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75
- Reaktionsverhalten Alkinyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69
- Synthese, Kristallstruktur und Reaktionswege von 2,2-Dimethyl-1,1,1,3,3,3-hexaphenyltrisilan (U. Baumeister, K. Schenzel und K. Hassler), 93
- Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4-nitrene ligand. Crystal structures of $[Ru_6(CO)_{15}(\mu\text{-}CO)_2(\mu_4\text{-}NH)(\mu\text{-}OMe)(\mu_3\text{-}\eta^2\text{-}N(H)C(O)OMe)]$ and $[Ru_6(CO)_{16}(\mu\text{-}CO)_2(\mu_4\text{-}NH)(\mu\text{-}OMe)(\mu\text{-}NCO)]$ (K.K.H. Lee and W.T. Wong), C43
- Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_5H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29
- Synthesis and structure of the dimeric organo-antimony (V) imido complex $[Ph_3Sb(\mu\text{-}NCH_2CH_2Ph)]_2$ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29
- The cleavage of a coordinated SCNR group in $[Fe(CO)_2L_2(\eta^2\text{-}SCNR)]$ by $[Co(\eta\text{-}C_5H_5)(PPh_3)_2]$ to give complexes containing μ_3-CNR ligand. The preparation and structure of $[\{Co(\eta\text{-}C_5H_5)\}_2\{Fe(CO)_2(PPh_3)\}(\mu_3\text{-}S)(\mu_3\text{-}CNC(O)C_6H_5)]$ (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46
- The relationship between the molecular structure of [3,3-$(PMe_2Ph)_2$-closo-3,1,2-$PtC_2B_9H_{11}$] and the mechanism of the fluxional behaviour of $M(PR_3)_2$-units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12

C_2-symmetrical
- Crystal structure of enantiomerically pure bis(dimethylbicyclooctylcyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

Cyclometallation
- Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β-unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Cyclopentadienyl
Coordination of the (μ-η^2-As_2) ligand to Group 6 transition metal carbonyl fragments and crystal structures of $Cp_2Cr_2(CO)_4As_2$-$[M(CO)_5]_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47
Crystal structure of enantiomerically pure bis(dimethylbicyclooctylcyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53
Cyclopentadienylcobalt complexes of some azepines (H. Wadepohl and K. Töllner), 111
Intramolekulare Insertion eines η^5-Cyclopentadienyl-Ringes in einem Bis-η^5-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefeber, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221
Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117
Protonation of $Cp(CO)(PPh_3)RuC{\equiv}CPh$: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct $[Cp(CO)(PPh_3)Ru{=}\overbrace{CCH(Ph)CH_2CH_2O}]^+$ (P. Nombel, N. Lugan and R. Mathieu), C22
Cyclopentadienyl complexes
Umsetzungen der lithiierten Halbsandwich-Komplexe $(CO)_3MC_5H_4$-Li (M = Mn, Re) mit organischen Carbonylverbindungen (M. Herberhold und M. Biersack), 277
Cyclosilanes
Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129

Dehydrogenative coupling
The X-ray crystal structures of the *ansa*-metallocenes, Me_2-$C(C_5H_4)_2MCl_2$ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205
Desulfurization
Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-*S*-benzoylsulphenamide. Crystal and molecular structures of $[Os_6(CO)_{20}(\mu_4\text{-}S)(MeCN)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19
Difluorocarbene
Donor-free bis(trifluoromethyl)cadmium, $(CF_3)_2Cd$: a readily available low-temperature difluorocarbene source (R. Eujen and B. Hoge), C51
Dimer
Synthesis and structure of the dimeric organo-antimony (V) imido complex $[Ph_3Sb(\mu\text{-}NCH_2CH_2Ph)]_2$ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29
Dinitrogen
Synthesis and chacterization of low valent complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1
Dinuclear
Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)-β-diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal–organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101
Dinuclear complexes
Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of $[\{Pt(2,9\text{-dimethyl-}1,10\text{-phenanthroline})\text{-}(Z\text{-}MeO_2CCH{=}CH\text{-}CO_2Me)(H_2O)\}_2Hg](BF_4)_2$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251
Diphosphine
Procedure for the degradation of 1,2-$(PR_2)_2$-1,2-dicarba-*closo*-dodecaborane(12) and 1-(PR_2)-2-R'-1,2-dicarba-*closo*-dodecaborane(12) (F. Teixidor, C. Viñas, M.M. Abad, R. Nuñez, R. Kivekäs and R. Sillanpää), 193
Disubstituted permethylcyclohexasilanes
Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129

Early transition metals
Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_5H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29
Electrochemistry
Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243
Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259
Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiénayl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171
Electrophilic addition
Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic $[Mn_3(CO)_{12}(\mu\text{-}H)]^{2-}$ and $[Fe_3(CO)_{11}]^{2-}$ anions (O. Rossell, M. Seco and G. Segalés), 225
Enantioface binding
Crystal structures of diastereomeric styrene complexes of the chiral rhenium fragment $[(\eta^5\text{-}C_5H_5)Re(NO)(PPh_3)]^+$: effect of ligating enantioface upon bonding and conformation (M. Sanaú, T.-S. Peng, A.M. Arif and J.A. Gladysz), 235
Enone
Conjugate addition reactions of α-azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213
EPR Spectroscopy
Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243
Ethylene
The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15

Ferrocene
Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259
Ferrocenecarbaldehyde
Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259
Ferrocenes
Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiénayl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171
[3]Ferrocenophanes
Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium

(P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243

Fluxionality
Cyclopentadienylcobalt complexes of some azepines (H. Wadepohl and K. Töllner), 111
The relationship between the molecular structure of [3,3-$(PMe_2Ph)_2$-closo-3,1,2-$PtC_2B_9H_{11}$] and the mechanism of the fluxional behaviour of $M(PR_3)_2$-units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12

Glucose complexes
Transition metal carbohydrate chemistry. Part 4. Homoleptic diacetoneglucose complexes of molybdenum and tungsten: chiral ligands associated with an M≡M triple bond (U. Piarulli, D.N. Williams, C. Floriani, G. Gervasio and D. Viterbo), 185

Gold
Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic $[Mn_3(CO)_{12}(\mu\text{-}H)]^{2-}$ and $[Fe_3(CO)_{11}]^{2-}$ anions (O. Rossell, M. Seco and G. Segalés), 225

Group 15
Synthesis and structure of the dimeric organo-antimony (V) imido complex $[Ph_3Sb(\mu\text{-}NCH_2CH_2Ph)]_2$ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29

Hafnium
The X-ray crystal structures of the *ansa*-metallocenes, $Me_2C(C_5H_4)_2MCl_2$ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205

^{1}H NMR
An ^{1}H and ^{13}C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ^{13}C–^{11}B and ^{13}C–^{205}Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265

Hydrogen bonding
Hydrogen-bonding 'Pd(OH)...2(HOAr)' and '$[Pd(OH_2)^+ \ldots {}^-OAr]_2$', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts $[Pd(OH)(C_4H_8)((pz)_3BH).2(3\text{-}MeC_6H_4OH)]$ and $[Pd(OH_2)(C_4H_8)((pz)_3BH).(C_6F_5O)]_2$ [$(pz)_3BH$ = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16

Hydroxo-ligand
Hydrogen-bonding 'Pd(OH)...2(HOAr)' and '$[Pd(OH_2)^+ \ldots {}^-OAr]_2$', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts $[Pd(OH)(C_4H_8)((pz)_3BH).2(3\text{-}MeC_6H_4OH)]$ and $[Pd(OH_2)(C_4H_8)((pz)_3BH).(C_6F_5O)]_2$ [$(pz)_3BH$ = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16

Imidazole
Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplexkationen: Synthese und Strukturen von $[R_3Sn(L)(L')]^+[(MeSO_2)_2N]^-$ (L = L' = HMPA, Ph_3AsO, Imidazol, 4-Dimethylaminopyridin; L = H_2O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155

Imido
Synthesis and structure of the dimeric organo-antimony (V) imido complex $[Ph_3Sb(\mu\text{-}NCH_2CH_2Ph)]_2$ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29

Intramolecular cyclization
A new tetradentate nitrogen ligand for organometallic catalysis: electrochemical nickel-catalyzed intramolecular cyclization (J.C. Clinet and E. Duñach), C48

Iridium
Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243

Iron
Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic $[Mn_3(CO)_{12}(\mu\text{-}H)]^{2-}$ and $[Fe_3(CO)_{11}]^{2-}$ anions (O. Rossell, M. Seco and G. Segalés), 225
Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117
Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259
The cleavage of a coordinated SCNR group in $[Fe(CO)_2L_2(\eta^2\text{-}SCNR)]$ by $[Co(\eta\text{-}C_5H_5)(PPh_3)_2]$ to give complexes containing μ_3-CNR ligand. The preparation and structure of $[\{Co(\eta\text{-}C_5H_5)\}_2\{Fe(CO)_2(PPh_3)\}(\mu_3\text{-}S)(\mu_3\text{-}CNC(O)C_6H_5)]$ (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46
Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiényl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171

μ_3-Isocyanide
The cleavage of a coordinated SCNR group in $[Fe(CO)_2L_2(\eta^2\text{-}SCNR)]$ by $[Co(\eta\text{-}C_5H_5)(PPh_3)_2]$ to give complexes containing μ_3-CNR ligand. The preparation and structure of $[\{Co(\eta\text{-}C_5H_5)\}_2\{Fe(CO)_2(PPh_3)\}(\mu_3\text{-}S)(\mu_3\text{-}CNC(O)C_6H_5)]$ (A.R. Manning, L. O'Dwyer, P.A. McArdle and D. Cunningham), C46

Isocyanide complexes
Formation of a rhenium(III) carbonyl complex by electrophilic attack on rhenium isocyanides. Synthesis and molecular structure of $\{Re[N(CH_2CH_2S)_3][CNC(CH_3)_3]\}$ and $\{Re[N(CH_2CH_2S)_3](CO)\}$ (M. Glaser, H. Spies, T. Lügger and F.E. Hahn), C32

Isomerisan
Protonation of $Cp(CO)(PPh_3)RuC{\equiv}CPh$: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct $[Cp(CO)(PPh_3)Ru{=}\overline{CCH(Ph)CH_2CH_2O}]^+$ (P. Nombel, N. Lugan and R. Mathieu), C22

Lithiated half-sandwich complexes
Umsetzungen der lithiierten Halbsandwich-Komplexe $(CO)_3MC_5H_4Li$ (M = Mn, Re) mit organischen Carbonylverbindungen (M. Herberhold und M. Biersack), 277

Lithium
XIII. Synthese und Eigenschaften von $[Li(TMEDA)]_2[Pt(CH_2CMe_2CMe_2CH_2)_2]$ (1); Vergleich der Zersetzungstemperaturen von Metallacyclopentan-Komplexen des Pt(II) und Ni(II) des Typs (1) mit und ohne β-Wasserstoff; Kristallstruktur von $(COD)PtCH_2SiMe_2SiMe_2CH_2$ (8) (R. Wyrwa, H.-O. Fröhlich und H. Görls), 135

Manganese
Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic $[Mn_3(CO)_{12}(\mu\text{-}H)]^{2-}$ and $[Fe_3(CO)_{11}]^{2-}$ anions (O. Rossell, M. Seco and G. Segalés), 225

Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β-unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59

Umsetzungen der lithiierten Halbsandwich-Komplexe $(CO)_3MC_5H_4$-Li (M = Mn, Re) mit organischen Carbonylverbindungen (M. Herberhold und M. Biersack), 277

Mechanism

The relationship between the molecular structure of $[3,3\text{-}(PMe_2Ph)_2\text{-}closo\text{-}3,1,2\text{-}PtC_2B_9H_{11}]$ and the mechanism of the fluxional behaviour of $M(PR_3)_2$-units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12

Mercury

Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of $[\{Pt(2,9\text{-dimethyl-}1,10\text{-phenanthroline})\text{-}(Z\text{-}MeO_2CCH{=}CHCO_2Me)(H_2O)\}_2Hg](BF_4)_2$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251

Metallacarborane

The relationship between the molecular structure of $[3,3\text{-}(PMe_2Ph)_2\text{-}closo\text{-}3,1,2\text{-}PtC_2B_9H_{11}]$ and the mechanism of the fluxional behaviour of $M(PR_3)_2$-units in closo-twelve-atom metallaheteroboranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12

Metallacycle

Hydrogen-bonding 'Pd(OH)...2(HOAr)' and '$[Pd(OH_2)^+ \ldots {}^-OAr]_2$', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts $[Pd(OH)(C_4H_8)\{(pz)_3BH\}.2(3\text{-}MeC_6H_4OH)]$ and $[Pd(OH_2)(C_4H_8)\{(pz)_3BH\}.(C_6F_5O)]_2$ [$(pz)_3BH$ = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16

Intramolekulare Insertion eines η^3-Cyclopentadienyl-Ringes in einem Bis-η^5-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefeber, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221

Preparation of cyclomanganated chalcones and their reactions with methyl acrylate and other α,β-unsaturated carbonyl compounds (W. Tully, L. Main and B.K. Nicholson), 75

Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289

Metallacyclopentanes

XIII. Synthese und Eigenschaften von $[Li(TMEDA)]_2[Pt(CH_2CMe_2CMe_2CH_2)_2]$ (1); Vergleich der Zersetzungstemperaturen von Metallacyclopentan-Komplexen des Pt(II) und Ni(II) des Typs (1) mit und ohne β-Wasserstoff; Kristallstruktur von $(COD)Pt(CH_2SiMe_2SiMe_2CH_2)$ (8) (R. Wyrwa, H.-O. Fröhlich und H. Görls), 135

Metallation

Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C–Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149

Synthesis and structure of the dimeric organo-antimony (V) imido complex $[Ph_2Sb(\mu\text{-}NCH_2CH_2Ph)]_2$ (A.J. Edwards, M.A. Paver, P. Pearson, P.R. Raithby, M.-A. Rennie, C.A. Russell and D.S. Wright), C29

Metallocene

Crystal structure of enantiomerically pure bis(dimethylbicyclooctylcyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

ansa-Metallocenes

The X-ray crystal structures of the *ansa*-metallocenes, $Me_2C(C_5H_4)_2MCl_2$ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205

Metal–metal bonds

Electrophilic additions of mono-, di- and tri-gold units to the isoelectronic $[Mn_3(CO)_{12}(\mu\text{-}H)]^{2-}$ and $[Fe_3(CO)_{11}]^{2-}$ anions (O. Rossell, M. Seco and G. Segalés), 225

Metal–sugar compounds

Transition metal carbohydrate chemistry. Part 4. Homoleptic diacetoneglucose complexes of molybdenum and tungsten: chiral ligands associated with an M≡M triple bond (U. Piarulli, D.N. Williams, C. Floriani, G. Gervasio and D. Viterbo), 185

Methoxide

Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26

Methoxycarbonyl complexes

Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26

MOCVD

Metallorganische Verbindungen des Kupfers. IX. Ein- und zweikernige Kupfer(I)-β-diketonat-Komplexe mit schwefel-verbrückten Bisalkinen: Potentielle Precursorsysteme für die "metal–organic chemical vapor deposition" (G. Schmidt und U. Behrens), 101

Molybdenum

Synthesis and chacterization of low valent complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1

Monosubstituted bi(cyclohexasilanyl) derivatives

Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129

Mössbauer

Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of $[SnCl_3\{HB(4\text{-}Cl\text{-}3,5\text{-}Me_2Pz)_3\}]$ and $[PhSnCl_2\{HB(4\text{-}Cl\text{-}3,5\text{-}Me_2Pz)_3\}]$ (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297

Nickel catalysis

A new tetradentate nitrogen ligand for organometallic catalysis: electrochemical nickel-catalyzed intramolecular cyclization (J.C. Clinet and E. Duñach), C48

Nitrene

Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4-nitrene ligand. Crystal structures of $[Ru_6(CO)_{15}(\mu\text{-}CO)_2(\mu_4\text{-}NH)(\mu\text{-}OMe)\{\mu_3\text{-}\eta^2\text{-}N(H)C(O)OMe\}]$ and $[Ru_6(CO)_{16}(\mu\text{-}CO)_2(\mu_4\text{-}NH)(\mu\text{-}OMe)(\mu\text{-}NCO)]$ (K.K.H. Lee and W.T. Wong), C43

Nitrile

The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15

Nitrogen ligand

A new tetradentate nitrogen ligand for organometallic catalysis: electrochemical nickel-catalyzed intramolecular cyclization (J.C. Clinet and E. Duñach), C48

NMR spectroscopy

Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of $[SnCl_3\{HB(4\text{-}Cl\text{-}3,5\text{-}Me_2Pz)_3\}]$ and $[PhSnCl_2\{HB(4\text{-}Cl\text{-}3,5\text{-}Me_2Pz)_3\}]$ (G. Gioia Lob-

bia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297

Nuclear magnetic resonance
Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289

Open metallocene
Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_5H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29

Organocopper
Conjugate addition reactions of α-azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

Organocuprate
Conjugate addition reactions of α-azoalkylcuprate reagents (C.W. Alexander, S.-Y. Lin and R.K. Dieter), 213

Osmium
Metal cluster-induced desulphurization of *N*-(*p*-methoxybenzoyl)-*S*-benzoylsulphenamide. Crystal and molecular structures of $[Os_6(CO)_{20}(\mu_4\text{-}S)(MeCN)]$ (S.-M. Lee, K.-K. Cheung and W.-T. Wong), C19

Oxidation
Utilisation du cyanure de tétraéthylammonium pour la préparation de complexes cyclopentadiényl fer cyclohexadiényl cyanés. Etude de leur oxydation électrochimique et chimique (N. Guennec et C. Moinet), 171

Oxidative addition
Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C–Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149

Oxidative carbonylation
Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21

Palladium
Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26
Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21
Formation of P,C-chelated palladium complexes by phosphine-assisted oxidative addition of an aliphatic C–Cl bond (M. Portnoy, Y. Ben-David and D. Milstein), 149
Hydrogen-bonding 'Pd(OH) . . . 2(HOAr)' and '$[Pd(OH_2)^+ \ldots {}^-OAr]_2$', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts $[Pd(OH)(C_4H_8)((pz)_3BH).2(3\text{-}MeC_6H_4OH)]$ and $[Pd(OH_2)(C_4H_8)((pz)_3BH).(C_6F_5O)]_2$ [$(pz)_3BH$ = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16

Pentadienyl
Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_5H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29

Perfluoroalkoxide
Carbonylation of diamino-ligated methylpalladium(II) methoxide complexes (G.M. Kapteijn, M.J. Verhoef, M.A.F.H. Van den Broek, D.M. Grove and G. Van Koten), C26

Phenyl
Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129

Phosphanes
Reaktionsverhalten Alkinyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69

Phosphazene
Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59

Phosphine
Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117
Procedure for the degradation of 1,2-$(PR_2)_2$-1,2-dicarba-*closo*-dodecaborane(12) and 1-(PR_2)-2-R′-1,2-dicarba-*closo*-dodecaborane(12) (F. Teixidor, C. Viñas, M.M. Abad, R. Nuñez, R. Kivekäs and R. Sillanpää), 193
Synthesis and chacterization of low valent complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1

Phosphine ligand
Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, $[WI_2(CO)(\eta^2\text{-triphos})(\eta^2\text{-}MeC_2\text{-}Me)].0.75CH_2Cl_2$ (triphos = $PhP(CH_2CH_2PPh_2)_2$) (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8

Phosphines
Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_5H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29

Phosphinites
Reaktionsverhalten Alkinyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69

Phosphite
Synthesis and chacterization of low valent complexes of molybdenum containing the tridentate ligand $PhP(CH_2CH_2PPh_2)_2$ (TRI) (T.A. George and H.H. Hammund), C1

Phosphorus
Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplexkationen: Synthese und Strukturen von $[R_3Sn(L)(L')]^+[(MeSO_2)_2N]^-$ (L = L′ = HMPA, Ph_3AsO, Imidazol, 4-Dimethylaminopyridin; L = H_2O und L′ = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155

Photochemical synthesis
The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15

Planar chirality
New chiral chelating phosphine complexes containing tricarbonyl(η^6-arene)chromium for highly enantioselective allylic alkylation (Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura), 143

Platinum
Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of [[Pt(2,9-dimethyl-1,10-phenanthroline)-(Z-$MeO_2CCH{=}CHCO_2Me$)(H_2O)]$_2$Hg]$(BF_4)_2$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251
The relationship between the molecular structure of [3,3-$(PMe_2Ph)_2$-closo-3,1,2-$PtC_2B_9H_{11}$] and the mechanism of the fluxional behaviour of $M(PR_3)_2$-units in closo-twelve-atom metallaheterobo-

ranes (D. O'Connell, T.R. Spalding, G. Ferguson, J.F. Gallagher and J.D. Kennedy), C12

XIII. Synthese und Eigenschaften von $[Li(TMEDA)]_2$-$Pt(CH_2CMe_2CMe_2CH_2)_2$ (1); Vergleich der Zersetzungstemperaturen von Metallacyclopentan-Komplexen des Pt(II) und Ni(II) des Typs (1) mit und ohne β-Wasserstoff; Kristallstruktur von $(COD)PtCH_2SiMe_2SiMe_2CH_2$ (8) (R. Wyrwa, H.-O. Fröhlich und H. Görls), 135

Polymeric

Synthesis and electrochemistry of polymeric complexes with cationic manganese carbonyl fragments coordinated to polyphosphazenes through nitrile ligands (G.A. Carriedo, L. Fernández-Catuxo, F.J.G. Alonso and P. Gómez-Elipe), 59

Potassium salts

An ^1H and ^{13}C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ^{13}C–^{11}B and ^{13}C–^{205}Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265

Pyrazolylborates

An ^1H and ^{13}C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ^{13}C–^{11}B and ^{13}C–^{205}Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265

Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of $[SnCl_3\{HB(4\text{-}Cl\text{-}3,5\text{-}Me_2Pz)_3\}]$ and $[PhSnCl_2\{HB(4\text{-}Cl\text{-}3,5\text{-}Me_2Pz)_3\}]$ (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297

Pyridine-*N*-oxide

Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplexkationen: Synthese und Strukturen von $[R_3Sn(L)(L')]^+[(MeSO_2)_2N]^-$ (L = L' = HMPA, Ph_3AsO, Imidazol, 4-Dimethylaminopyridin; L = H_2O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155

Reductive carbonylation

Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21

Rhenium

Crystal structures of diastereomeric styrene complexes of the chiral rhenium fragment $[(\eta^5\text{-}C_5H_5)Re(NO)(PPh_3)]^+$: effect of ligating enantioface upon bonding and conformation (M. Sanaú, T.-S. Peng, A.M. Arif and J.A. Gladysz), 235

Formation of a rhenium(III) carbonyl complex by electrophilic attack on rhenium isocyanides. Synthesis and molecular structure of $\{Re[N(CH_2CH_2S)_3][CNC(CH_3)_3]\}$ and $\{Re[N(CH_2CH_2S)_3](CO)\}$ (M. Glaser, H. Spies, T. Lügger and F.E. Hahn), C32

Umsetzungen der lithiierten Halbsandwich-Komplexe $(CO)_3MC_5H_4$-Li (M = Mn, Re) mit organischen Carbonylverbindungen (M. Herberhold und M. Biersack), 277

Rhodium

Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243

Ruthenium

Protonation of $Cp(CO)(PPh_3)RuC{\equiv}CPh$: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct $[Cp(CO)(PPh_3)Ru{=}CCH(Ph)CH_2CH_2O]^+$ (P. Nombel, N. Lugan and R. Mathieu), C22

Synthesis and characterisation of hexanuclear ruthenium clusters containing a μ_4-nitrene ligand. Crystal structures of $[Ru_6(CO)_{15}(\mu\text{-}CO)_2(\mu_4\text{-}NH)(\mu\text{-}OMe)\{\mu_3\text{-}\eta^2\text{-}N(H)C(O)OMe\}]$ and $[Ru_6(CO)_{16}(\mu\text{-}CO)_2(\mu_4\text{-}NH)(\mu\text{-}OMe)(\mu\text{-}NCO)]$ (K.K.H. Lee and W.T. Wong), C43

The photochemical generation of novel neutral mononuclear ruthenium complexes and their reactivity (A.J. Edwards, N.E. Leadbeater, J. Lewis and P.R. Raithby), 15

Silicon

Synthese, Kristallstruktur und Reaktionswege von 2,2-Dimethyl-1,1,1,3,3,3-hexaphenyltrisilan (U. Baumeister, K. Schenzel und K. Hassler), 93

Silver

Reaktionsverhalten Alkinyl-substituierter Phosphinite und Phosphane am Beispiel von Kupfer(I)- und Silber(I)-Trifluormethansulfonaten (H. Lang, M. Weinmann, M. Winter, M. Leise und W. Imhof), 69

^{29}Si-NMR spectroscopy

Synthese, Kristallstruktur und Reaktionswege von 2,2-Dimethyl-1,1,1,3,3,3-hexaphenyltrisilan (U. Baumeister, K. Schenzel und K. Hassler), 93

Stannyl

Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289

Styrene complexes

Crystal structures of diastereomeric styrene complexes of the chiral rhenium fragment $[(\eta^5\text{-}C_5H_5)Re(NO)(PPh_3)]^+$: effect of ligating enantioface upon bonding and conformation (M. Sanaú, T.-S. Peng, A.M. Arif and J.A. Gladysz), 235

Sulfonyl

Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplexkationen: Synthese und Strukturen von $[R_3Sn(L)(L')]^+[(MeSO_2)_2N]^-$ (L = L' = HMPA, Ph_3AsO, Imidazol, 4-Dimethylaminopyridin; L = H_2O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155

Tetradentate

A new tetradentate nitrogen ligand for organometallic catalysis: electrochemical nickel-catalyzed intramolecular cyclization (J.C. Clinet and E. Duñach), C48

Thallium salts

An ^1H and ^{13}C NMR spectroscopic study of the structure of potassium and thallium salts of tris- and tetrakis-(pyrazol-1-yl) borates in solution. Some ^{13}C–^{11}B and ^{13}C–^{205}Tl residual coupling constants (C. López, D. Sanz, R.M. Claramunt, S. Trofimenko and J. Elguero), 265

Theoretical calculations

Electrochemical, spectroscopic and theoretical studies of a series of 1,1'-ferrocene dichalcogenido-complexes of rhodium and iridium (P. Zanello, M. Casarin, L. Pardi, M. Herberhold and G.-X. Jin), 243

Thiourea

Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide–thiourea catalysts (B. Gabriele, G. Salerno, M. Costa and G.P. Chiusoli), 21

Tin

Organotin (IV) polypyrazolyborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of $[SnCl_3\{HB(4\text{-}Cl\text{-}3,5\text{-}$

$Me_2Pz)_3$] and [$PhSnCl_2${HB(4-Cl-3,5-Me_2Pz)}$_3$] (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297

Polysulfonylamine LXII. Dimesylamid als nichtkoordinierendes Gegenion für Organozinn-Komplexkationen: Synthese und Strukturen von [$R_3Sn(L)(L')$]$^+$[$(MeSO_2)_2N$]$^-$ (L = L' = HMPA, Ph_3AsO, Imidazol, 4-Dimethylaminopyridin; L = H_2O und L' = Pyridin-*N*-oxid) (I. Lange, D. Henschel, A. Wirth, J. Krahl, A. Blaschette und P.G. Jones), 155

Stepwise synthesis of organometallic-substituted stannoles via stabilized triorganotin cations (B. Wrackmeyer, K. Horchler von Locquenghien and S. Kundler), 289

Tetradentate -N_2O_2 ligand complexes of Tin(II). X-ray crystal structure of [*N,N'*-(1,2-ethylene)*bis*(salicylaldamine)]tin(II), ($SaleanH_2Sn$) (D.A. Atwood, J.A. Jegier, K.J. Martin and D. Rutherford), C4

Titanium

The X-ray crystal structures of the *ansa*-metallocenes, $Me_2C(C_5H_4)_2MCl_2$ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205

Titanocene

Intramolekulare Insertion eines η^5-Cyclopentadienyl-Ringes in einem Bis-η^5-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefeber, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221

Transmetallation

Tetradentate -N_2O_2 ligand complexes of Tin(II). X-ray crystal structure of [*N,N'*-(1,2-ethylene)*bis*(salicylaldamine)]tin(II), ($SaleanH_2Sn$) (D.A. Atwood, J.A. Jegier, K.J. Martin and D. Rutherford), C4

Triflate

Synthesis of bi(cyclohexasilanyl) derivatives (P. Gspaltl, A. Spielberger, A. Zechmann and E. Hengge), 129

Trifluoromethyl

Donor-free bis(trifluoromethyl)cadmium, $(CF_3)_2Cd$: a readily available low-temperature difluorocarbene source (R. Eujen and B. Hoge), C51

Trimethylsilyl

Intramolekulare Insertion eines η^5-Cyclopentadienyl-Ringes in einem Bis-η^5-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefeber, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221

Trinuclear complexes

Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of [{Pt(2,9-dimethyl-1,10-phenanthroline)-(Z-$MeO_2CCH{=}CH$-CO_2Me)(H_2O)}$_2$Hg](BF_4)$_2$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251

Triphos, $PhP(CH_2CH_2PPh_2)_2$

Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, [$WI_2(CO)(\eta^2$-triphos$)(\eta^2$-MeC_2-Me)].0.75CH_2Cl_2 {triphos = $PhP(CH_2CH_2PPh_2)_2$} (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8

Trisilanes

Synthese, Kristallstruktur und Reaktionswege von 2,2-Dimethyl-1,1,1,3,3,3-hexaphenyltrisilan (U. Baumeister, K. Schenzel und K. Hassler), 93

Tungsten

Coordination of the (μ-η^2-As_2) ligand to Group 6 transition metal carbonyl fragments and crystal structures of $Cp_2Cr_2(CO)_4As_2$-[$M(CO)_5$]$_2$ (M = Cr, W) (L.Y. Goh, W. Chen and R.C.S. Wong), 47

Tungsten(II)

Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, [$WI_2(CO)(\eta^2$-triphos$)(\eta^2$-MeC_2-Me)].0.75CH_2Cl_2 {triphos = $PhP(CH_2CH_2PPh_2)_2$} (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8

Vinyl

Preparations of novel half-sandwich iron carbonyl phosphine complexes: intramolecular link between the cyclopentadienyl and the phosphine ligand (T.-F. Wang, J.-P. Juang and Y.-S. Wen), 117

Vinylidene complex

Protonation of $Cp(CO)(PPh_3)RuC{\equiv}CPh$: Formation of cationic phenylacetylene and phenylvinylidene complexes and subsequent reaction with ethylene oxide to form the net [2 + 3] cycloadduct [$Cp(CO)(PPh_3)Ru{=}\overbrace{CH(Ph)CH_2CH_2O}$]$^+$ (P. Nombel, N. Lugan and R. Mathieu), C22

Water

Hydrogen-bonding 'Pd(OH) . . . 2(HOAr)' and '[$Pd(OH_2)$]$^+$ . . . $^-$OAr]$_2$', and protonation of a hydroxo-ligand by pentafluorophenol in organopalladium(IV) chemistry. Structural studies of the pallada(IV)cyclopentane adducts [Pd(OH)(C_4H_8)($pz)_3$BH].2(3-MeC_6H_4OH)] and [$Pd(OH_2)(C_4H_8)$($pz)_3$BH].(C_6F_5O)$_2$ [($pz)_3$BH = tris(pyrazol-1-yl)borate] (A.J. Canty, H. Jin, B.W. Skelton and A.H. White), C16

X-ray analysis

Bi- and trinuclear cationic complexes involving bonds between mercury and five-coordinate platinum(II). Molecular structure of [{Pt(2,9-dimethyl-1,10-phenanthroline)-(Z-$MeO_2CCH{=}CH$-CO_2Me)(H_2O)}$_2$Hg](BF_4)$_2$ (M.E. Cucciolito, F. Giordano, F. Ruffo and V. De Felice), 251

X-ray crystallography

Formation of a rhenium(III) carbonyl complex by electrophilic attack on rhenium isocyanides. Synthesis and molecular structure of {Re[$N(CH_2CH_2S)_3$][$CNC(CH_3)_3$]} and {Re[$N(CH_2CH_2S)_3$](CO)} (M. Glaser, H. Spies, T. Lügger and F.E. Hahn), C32

X-ray diffraction

Intramolekulare Insertion eines η^5-Cyclopentadienyl-Ringes in einem Bis-η^5-cyclopentadienyltitanacyclopentadien (U. Rosenthal, C. Lefeber, P. Arndt, A. Tillack, W. Baumann, R. Kempe und V.V. Burlakov), 221

X-ray structure

Organotin (IV) polypyrazolylborates. VIII. Hydridotris (4-chloro-3,5-dimethyl-1*H*-pyrazol-1-yl)borates. Characterization, Mössbauer study and X-ray crystal structures of [$SnCl_3${HB(4-Cl-3,5-$Me_2Pz)_3$}] and [$PhSnCl_2${HB(4-Cl-3,5-$Me_2Pz)_3$}] (G. Gioia Lobbia, P. Cecchi, S. Calogero, G. Valle, M. Chiarini and L. Stievano), 297

Reaction of ferrocenecarbaldehyde with *o*-phenylenediamine. Crystal structure of *N*-ferrocenylmethyl-2-ferrocenyl-benzimidazole (A. Benito, R. Martínez-Máñez, J. Payá, J. Soto, M.J.L. Tendero and E. Sinn), 259

Synthesis, molecular structure and reactions of the novel organometallic phosphine ligand, [$WI_2(CO)(\eta^2$-triphos$)(\eta^2$-MeC_2-Me)].0.75CH_2Cl_2 {triphos = $PhP(CH_2CH_2PPh_2)_2$} (P.K. Baker, S.J. Coles, M.B. Hursthouse, M.M. Meehan and S.D. Ridyard), C8

XIII. Synthese und Eigenschaften von [Li(TMEDA)]$_2$-$\overbrace{Pt(CH_2CMe_2CMe_2CH_2)_2}$ (1); Vergleich der Zersetzungstemperaturen von Metallacyclopentan-Komplexen des Pt(II) und Ni(II)

des Typs (1) mit und ohne β-Wasserstoff; Kristallstruktur von (COD)PtCH$_2$SiMe$_2$SiMe$_2$CH$_2$ (8) (R. Wyrwa, H.-O. Fröhlich und H. Görls), 135

Ytterbium
Synthesis and structure of a new organolanthanide oxide complex, $(CH_3C_5H_4)_3Yb_4(\mu\text{-Cl})_6(\mu_3\text{-Cl})(\mu_4\text{-O})(THF)_3$ (X. Zhou, H. Ma, Z. Wu, X. You, Z. Xu and X. Huang), 11

Ziegler–Natta polymerization
Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307

Zirconium
Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307
Synthesis and characterization of phosphine adducts of the open zirconocene $Zr(C_3H_7)_2$ (T.E. Waldman, A.L. Rheingold and R.D. Ernst), 29
The X-ray crystal structures of the *ansa*-metallocenes, $Me_2C(C_5H_4)_2MCl_2$ (M = Ti, Zr and Hf) (R.M. Shaltout, J.Y. Corey and N.P. Rath), 205

Zirconocene
Bis(trimethylsilyl)benzamidinate zirconium dichlorides. Active catalysts for ethylene polymerization (D. Herskovics-Korine and M.S. Eisen), 307
Crystal structure of enantiomerically pure bis(dimethylbicyclooctylcyclopentadienyl)zirconium dichloride (R.L. Halterman and Z. Chen), 53

Book reviews
Advances in Silicon Chemistry Vol. 2. G.L. Larson (Ed.) (C. Eaborn), C36
Dictionary of organometallic compounds. Second Edition. B.J. Aylett, M.F. Lappert and P.L. Pauson (Eds.) (C. Eaborn), C36
Fieser and Fieser's reagents for organic synthesis, M. Fieser (C. Eaborn), C38
Gmelin Handbook of Inorganic and Organometallic Chemistry. 8th Edition *Organogermanium compounds*. Part 4. *Compounds with germanium–hydrogen bonds*, U. Krüerke, C. Siebert and B. Wöbke (Eds.) (C. Eaborn), C37
Gmelin Handbook of Inorganic and Organometallic Chemistry. 8th Edition *Sn organotin compounds*. Part 21. U. Krüerke (Ed.) (C. Eaborn), C37
Gmelin Handbook of Inorganic and Organometallic Chemistry, 8th Edition, Au, Compounds with Br, I, S, Se, Te, Po, B, and C, Supplement volume B2 G.J. Leigh (R. Eajen), C55
Organic Reactions. Equilibria, Kinetics and Mechanism, F. Ruff and I.G. Csizmadia (C. Eaborn), C37